

CRH LIBRARY

UNITED STATES
DEPARTMENT OF
COMMERCE
PUBLICATION



NOAA Technical Memorandum NWS WR70

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Weather Service

Predicting Inversion Depths and Temperature Influences in the Helena Valley

DAVID E. OLSEN

Western Region

SALT LAKE CITY,
UTAH

October 1971

NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 23 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS.

Papers 1 to 23, except for 5 (revised edition) and 10, are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84111. Papers 5 (revised edition), 10, and all others beginning with 24 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Bldg., 5285 Port Royal Road, Springfield, Va. 22151. Price: \$3.00 paper copy; \$0.95 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda

- WRTM 1 Some Notes on Probability Forecasting. Edward D. Diemer, September 1965. (Out of print.)
WRTM 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
WRTM 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965 to February 20, 1966. Edward D. Diemer, March 1966.
WRTM 4 Use of Meteorological Satellite Data. March 1966.
WRTM 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-178000)
WRTM 6 Improvement of Forecast Wording and Format. C. L. Glenn, May 1966.
WRTM 7 Final Report on Precipitation Probability Test Programs. Edward D. Diemer, May 1966.
WRTM 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967). (Out of print.)
WRTM 9 A Collection of Papers Related to the 1966 NMC Primitive-Equation Model. June 1966.
WRTM 10 Sonic Boom. Loren Crow (6th Weather Wing, USAF, Pamphlet), June 1966. (Out of print.) (AD-479366)
WRTM 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
WRTM 12 A Comparison of Fog Incidence at Missoula, Montana, with Surrounding Locations. Richard A. Dightman, August 1966. (Out of print.)
WRTM 13 A Collection of Technical Attachments on the 1966 NMC Primitive-Equation Model. Leonard W. Snellman, August 1966. (Out of print.)
WRTM 14 Application of Net Radiometer Measurements to Short-Range Fog and Stratus Forecasting at Los Angeles. Frederick Thomas, September 1966.
WRTM 15 The Use of the Mean as an Estimate of "Normal" Precipitation in an Arid Region. Paul C. Kangieser, November 1966.
WRTM 16 Some Notes on Acclimatization in Man. Edited by Leonard W. Snellman, November 1966.
WRTM 17 A Digitalized Summary of Radar Echoes Within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
WRTM 18 Limitations of Selected Meteorological Data. December 1966.
WRTM 19 A Grid Method for Estimating Precipitation Amounts by Using the WSR-57 Radar. R. Granger, December 1966. (Out of print.)
WRTM 20 Transmitting Radar Echo Locations to Local Fire Control Agencies for Lightning Fire Detection. Robert R. Peterson, March 1967. (Out of print.)
WRTM 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
WRTM 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.
WRTM 23 "K" Chart Applications to Thunderstorm Forecasts Over the Western United States. Richard E. Hambidge, May 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- WBTM 24 Historical and Climatological Study of Grinnell Glacier, Montana. Richard A. Dightman, July 1967. (PB-178071)
WBTM 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
WBTM 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
WBTM 27 Objective Minimum Temperature Forecasting for Helena, Montana. D. E. Olsen, February 1968. (PB-177827)
WBTM 28 Weather Extremes. R. J. Schmidli, April 1968 (revised July 1968). (PB-178928)
WBTM 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)
WBTM 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)
WBTM 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB-179084)
WBTM 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)
WBTM 33 Objective Forecasting. Philip Williams, Jr., August 1968. (AD-680425)
WBTM 34 The WSR-57 Radar Program at Missoula, Montana. R. Granger, October 1968. (PB-180292)
WBTM 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and DeVon B. Smith, December 1968 (revised June 1970). (AD-681857)
WBTM 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (Out of print.) (PB-183055)
WBTM 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)
WBTM 38 Climate of Phoenix, Arizona. R. J. Schmidli, P. C. Kangieser, and R. S. Ingram, April 1969. (Out of print.) (PB-184295)
WBTM 39 Upper-Air Lows Over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)
WBTM 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L. W. Snellman, August 1969. (PB-185068)
WBTM 41 High Resolution Radiosonde Observations. W. S. Johnson, August 1969. (PB-185673)
WBTM 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)
WBTM 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)
WBTM 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)
WBTM 45/1 Precipitation Probabilities in the Western Region Associated with Winter 500-mb Map Types. Richard A. Augulis, December 1969. (PB-188248)

U. S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL WEATHER SERVICE

NOAA Technical Memorandum NWSTM WR-70

PREDICTING INVERSION DEPTHS AND TEMPERATURE
INFLUENCES IN THE HELENA VALLEY

David E. Olsen
Weather Service Forecast Office
Boise, Idaho



WESTERN REGION
TECHNICAL MEMORANDUM NO. 70

SALT LAKE CITY, UTAH
OCTOBER 1971

TABLE OF CONTENTS

	<u>Page</u>
List of Figures and Tables	iii
I. Introduction	i
II. Development of the Inversion Depth Formulae	1-4
III. Temperature Influences Under Inversions	4-6
IV. Verification of Temperature Under Inversions	6-7
V. Conclusions	7-8
VI. Acknowledgments	8
VII. References	8

LIST OF FIGURES AND TABLES

	<u>Page</u>
Figure 1. Schematic Inversion and the Area Enclosed by the Inversion Curve and the Approximate Normal Lapse Rate Curve Extended from Height Z to the Ground to Temperature T_L .	9
Figure 2. Illustrating the Inversion Curve Divided into Two Unequal Areas in Order to Derive an Equation to Find the Total Area Enclosed by T_{MI} Z T_L .	10
Figure 3. Illustrating the Area Under the Inversion Curve Representing the Energy Necessary to Raise the Temperature from T_{MI} to T_{MA} , and the Division of this Area into Two Unequal Portions in Order to Derive an Equation to Evaluate the Area.	11
Figure 4. Relationship between A^{II} and $T_{MA}-T_{MI}$ for $a = 1.15$ and $a = 1.25$.	12
Figure 5. Four Examples Comparing the Actual Temperature Curve (Solid Line) with the Predicted Temperature Curve (Dashed Line).	13
Table 1. Comparison of Computed Depths of Morning Inversion Layers and Observed Depths for Selected Winter Soundings.	4
Table 2. Values of $T_{MA}-T_{MI}$ Corresponding to Selected Values of A^{II} for $a = 1.15$ (Snow Cover) and $a = 1.25$.	6
Table 3. Comparison of Predicted Temperatures Under Inversion Conditions and Observed Temperatures.	7

PREDICTING INVERSION DEPTHS AND TEMPERATURE INFLUENCES IN THE HELENA VALLEY

I. INTRODUCTION

Surface inversions in the Helena valley are common throughout the year with strongest and most frequent occurrence during summer and fall. The quantitative effect of inversions has been imperfectly understood because the depth and configuration have been unknown. The purpose of this paper is to present an objective method for predicting the depth of the local inversion and its influence on the maximum or hourly temperature using a remote temperature sounding (Great Falls, 90 miles away) and the local minimum temperature. This is accomplished by assuming the depth of the inversion under clear skies and calm winds is directly related to the minimum temperature and the "temperature" which would occur if no inversion formed.

II. DEVELOPMENT OF THE INVERSION DEPTH FORMULAE

From summer of 1970 to March 1971, the National Weather Service Office, Helena, received aircraft temperature soundings from surface to about 8000 feet MSL during mornings when it was suspected an inversion existed. However, no aircraft sounding was ever taken on Sunday (day of rest!). Soundings were obtained from a local flying service aircraft originally hired by American Smelting and Refinery of east Helena, and after September 1970, the National Health Service. Temperature sensors used were the aircraft's own thermometer prior to September 1970, and a thermometer loaned by Western Region Headquarters, National Weather Service, after September 1970.

Careful studies of some of the aircraft soundings (see Figure 5) indicate that local inversions often resemble the exponential curve.

$$(1) \quad Z = a^{\Delta T} - 1$$

where: Z = Equivalent height above the surface in C° units on an adiabatic chart.

ΔT = Temperature difference from the minimum (T_{\min}) to any temperature (T) in C° (T_{\min} will therefore always be the origin for equation (1)).

a = a constant.

Furthermore, the area (energy loss) under the inversion appears to be directly related to the difference between the surface temperature which would occur should no inversion exist, T_L , and the observed surface minimum T_{MI} . (See Figure 1.) In fact, 22 fall aircraft soundings considered ideal for the purpose revealed through linear regression techniques that:

$$(2) \quad A^I = 3.5 (T_L - T_{MI})$$

where: A^I = Area in C° square units on adiabatic chart D-11.

T_L = That temperature which would result if the sounding above the inversion were extended down to the surface.

T_{MI} = Observed minimum surface temperature.

Equation (2) is really the crux of the entire presentation. We know that an inversion forms by radiative and transfer processes; and since the surface temperature change from T_L to T_{MI} is a balance of these processes, it seems logical to relate the surface change to the entire inversion under clear skies and light winds. Also, the 22 ideal examples show the value of "a" in (1) to approximately equal 1.25. Figure 1 shows a typical clear night inversion. In order to evaluate the total area (A) as shown in Figure 1, we must divide it into two unequal parts as shown in Figure 2. First:

$$(3) \quad A'_1 = \int_0^{(T_0 - T_{mi})} (2^{\Delta T} - 1) d(\Delta T)$$

For A'_2 some definitions are made. The slope of any sounding extrapolated to the ground to T_L is assumed here to be such that some height Z will equal twice the base of the triangle ZT_LT_0 as shown in Figure 2. Therefore:

$$(4) \quad A'_2 = \frac{Z^2}{4}$$

Now the sum of the areas A'_1 and A'_2 can be expressed:

$$(5) \quad A' = \int_0^{(T_0 - T_{mi})} (2^{\Delta T} - 1) d(\Delta T) + \frac{Z^2}{4}$$

Integration of (5) gives:

$$(5a) \quad A' = \frac{a^{(T_o - T_{mi})} - 1}{\text{Loge } a} - (T_o - T_{mi}) + \frac{Z^2}{4}$$

Equation (5a) is still not in usable form. However, we know,

$$(6) \quad T_o = T_L - \frac{Z}{2}$$

$$(7) \quad Z = a^{(T_o - T_{mi})} - 1$$

By substituting (6) and (7) into (5a) we arrive at,

$$(8) \quad A' = \frac{Z}{\text{Loge } a} + \frac{Z}{2} + \frac{Z^2}{4} - (T_L - T_{mi})$$

By rearranging and substituting into the general quadratic formula,

$$(9) \quad Z = -2 \left(\frac{1}{\text{Loge } a} + \frac{1}{2} \right) + 2 \left[\left(\frac{1}{\text{Loge } a} + \frac{1}{2} \right)^2 + (T_L - T_{mi}) + A' \right]^{\frac{1}{2}}$$

Since "a" already has been said to equal 1.25 and A' has been found to equal 3.5 ($T_L - T_{mi}$), we can combine and substitute to obtain:

$$(10) \quad Z \approx -10 + 2 [25 + 4.5(T_L - T_{mi})]^{\frac{1}{2}}$$

Inversions recorded later in the season with snow cover indicate "a" in (1) to be close to 1.15, so during snow-cover conditions (10) becomes:

$$(11) \quad Z \approx -15 + 2 [58 + 4.5(T_L - T_{mi})]^{\frac{1}{2}}$$

Since Z is equal to C° units, to obtain feet one must multiply Z times 300 because on adiabatic plotting chart D-11 one centigrade degree equals about 300 feet.

Since all fall soundings appropriate to the problem were used to establish the constants and coefficients in the above equations, only winter soundings are at present available for testing equations (10) and (11). (It is hoped that the aircraft sounding program can again be carried on during the late summer and fall of 1971.) Most clear night inversions occur during summer and fall months. Also, soundings were never taken on Sundays. Nevertheless, some verification examples from some winter soundings are presented in Table I, despite lack of truly ideal conditions.

DATE	CHARACTER OF PAST 12 HRS.	OBSERVED DEPTH FEET	COMPUTED DEPTH FEET
11/17/70	P.C. windy most of night	600	600
11/18/70	M.C. windy at times with sprinkle of snow	700	1500
11/20/70	V.C. snow before midnight	1500	2100
12/02/70	V.C. light winds	1800	1600
12/03/70	M. clear	700	1000
12/05/70	High cloudiness	2700	2700
12/08/70	Cloudy, windy at times	200	300
12/10/70	Cloudy, trace snow	1600	1800
12/11/70	Cloudy	1500	1300
12/12/70	P.C. windy most of time	200	600
12/14/70	High and middle clouds	2300	2100

Table I. Comparison of computed depths of morning inversion layers and observed depths for selected winter soundings.

III. TEMPERATURE INFLUENCES UNDER INVERSIONS

Another approach can be developed which verifies (10) and (11) indirectly. If "a" truly equals 1.25 or 1.15 (snow cover), and if we select clear days when it is evident that ideal inversion conditions exist, we should be able to accurately predict a rise in temperature under the inversion because energy units in degree centigrade squares are known [1].

To do this we add a certain number of heating units under the inversion and compare computed and observed temperatures. To accomplish this, we must develop formulae.

Referring to Figure 3, the area A^{II} under the inversion curve, which represents the number of energy units necessary to raise the surface temperature from T_M to some temperature T_{MA} , is enclosed by the

inversion curve and a dry adiabat through T_{MA} . This area, A^{II} , can be divided into two portions. The first portion from T_{M1} to T is given by the integral:

$$\int_0^{(T-T_{M1})} (a^{\Delta T} - 1) d(\Delta T)$$

The remaining portion of A^{II} is enclosed by the triangle $ZT T_{MA}$. Since the slope of an adiabat is approximately 45° , the height Z of the triangle is approximately equal to the base and the area of the triangle can be expressed as:

$$\frac{Z^2}{2}$$

From (1)

$$Z = a^{\frac{(T-T_{M1})}{-1}} - 1$$

and

$$Z^2 = \left[a^{\frac{(T-T_{M1})}{-1}} - 1 \right]^2$$

Adding the two portions of A^{II} together:

$$A'' = \int_0^{(T-T_{M1})} (a^{\Delta T} - 1) d(\Delta T) + \frac{(a^{\frac{(T-T_{M1})}{-1}} - 1)^2}{2}$$

or

$$(12) \quad A'' = \frac{a^{\frac{(T-T_{M1})}{-1}} - 1}{\log a} - (T - T_{M1}) + \frac{(a^{\frac{(T-T_{M1})}{-1}} - 1)^2}{2}$$

By substituting values of T in (12) we, of course, get A^{II} units.

However, to find the temperature T_{MA} as shown in Figure 3 we must use the equation:

$$(13) \quad T_{MA} = T + a^{T-T_{MI}} - 1$$

Table 2 shows values of $T_{MA}-T_{MI}$ corresponding to selected values of A^{II} using "a" equaling both 1.15 and 1.25.

<u>A^{II}</u>	$a = 1.15$	$a = 1.25$
	<u>$T_{MA}-T_{MI}$</u>	<u>$T_{MA}-T_{MI}$</u>
2	4.5	4.0
3	6.0	5.0
4	7.0	6.0
6	8.5	7.0
10	10.0	9.0
12	11.5	9.5
16	13.0	10.5
21	14.5	12.0
27	16.0	13.0
35	18.0	14.5
46	20.0	16.0
59	22.0	17.5

Table 2. Values of $T_{MA}-T_{MI}$ corresponding to selected values of A^{II} for $a = 1.15$ (snow cover) and $a = 1.25$.

Curves showing the complete range of A^{II} relative to $T_{MA}-T_{MI}$ are plotted in Figure 4. By computing available energy units at noon from [1], we then proceed to the verification.

IV. VERIFICATION OF TEMPERATURE UNDER INVERSIONS

The verification procedure used in this instance is to attempt to predict the noon, or in three cases the maximum temperature at Helena when the station was under an inversion. Since energy units from sunrise until noon or maximum can be computed, it should be easy to predict a noon or maximum temperature provided "a" of the exponential curve is 1.15 or 1.25. Below, in Table 3, are a few examples under clear or cirrus days during 1969.

DATE	ENERGY U*	UNITS** USED	PREDICTED TEMPERATURE	OBSERVED TEMPERATURE	DIFF.
12/25/69	25	15	28	24	4
12/11/69	25	7	24	23	1
12/6/69	25	15	23	20	3
12/3/69	30	18	26	24	2
12/3/69	30	30	Max 30	33	-3
12/2/69	30	18	28	28	0
12/1/69	30	18	29	31	-2
12/1/69	30	30	Max 34	40	-6
11/30/69	30	18	31	29	+2
11/29/69	30	14	34	32	+2
11/29/69	30	25	Max 38	40	-2
11/28/69	30	18	32	30	2

Table 3. Comparison of predicted temperatures under inversion conditions and observed temperatures.

*Energy units available to reach maximum temperature on clear days.

**Estimated energy units needed to reach noon temperature (a maximum temperature in three cases) taking into consideration cloudiness and snow cover.

As Table 3 shows, predicted noon temperatures are, in general, within two or three degrees of observed values.

A more thorough verification has not been possible due to lack of suitable, clear night soundings. However, computations which have been made under close to ideal conditions of essentially clear skies and light winds result in an accuracy which leads to high confidence in the equations developed. Inversions forming under ideal conditions of clear skies and light winds may occur one out of every two or three nights in the Helena valley July through September, but only one out of every six nights the remainder of the year.

V. CONCLUSIONS

It has been shown that the temperature sounding in the very lowest levels on clear, relatively calm mornings in the Helena valley can be approximated rather accurately by an exponential curve. Formulae have been developed from which the depth of the early morning inversion layer may be computed as a function of the observed minimum temperature at Helena airport and a "temperature" which would have prevailed if no inversion had formed. This latter temperature is obtained by extending the upper portion of a nearby temperature sounding (Great Falls, Montana) downward to the surface. Verification of computed heights of inversions shows good accuracy, especially under ideal radiative conditions. For

a more thorough verification of the formulae developed in this study, additional summer and fall low-level aircraft soundings in the Helena valley are needed.

Computation of inversion depths for a number of clear nights during the past several summers and falls using the Great Falls sounding and the Helena minimum temperature indicates theoretically that inversion depths in the Helena valley of 1500 to 2000 feet are not uncommon and 3000 feet not too rare. Generally, inversions tend to be deeper in fall than any other time, which is not too surprising. In general, inversions in the Helena valley occur on the majority of all nights, and become strong on an estimated one hundred nights per year.

VI. ACKNOWLEDGMENTS

Appreciation is expressed to Mr. Woodrow W. Dickey, Scientific Services Division, Western Region Headquarters, for his helpful suggestions. Mr. Richard Dightman, Meteorologist in Charge, Weather Service Office, Helena, Montana, was primarily responsible for the arrangements with the National Health Service to continue the low-level aircraft sounding program.

VII. REFERENCES

1. Olsen, David E., "Forecasting Maximum Temperatures at Helena, Montana", Technical Memorandum WSTM WR-43, October 1969.
2. Jefferson, C. J., "Temperature Rise on Clear Mornings", The Meteorological Magazine, Vol. 79, No. 932, February 1950, page 33.




Figure 1: Schematic inversion and the area enclosed by the inversion curve and the approximate normal lapse rate curve extended from height Z to the ground to temperature T_L .




Figure 2. Illustrating the inversion curve divided into two unequal areas in order to derive an equation to find the total area enclosed by T_{M1} Z T_L .




Figure 3. Illustrating the area under the inversion curve representing the energy necessary to raise the temperature from T_{M1} to T_{MA} , and the division of this area into two unequal portions in order to derive an equation to evaluate the area.




Figure 4. Relationship between A'' and $TNA - TM_1 C^\circ$ for $a = 1.15$ and $a = 1.25$.




Figure 5. Four examples comparing the actual temperature curve (solid line) with the predicted temperature curve (dashed line).